

Transposition of a local-density-dependent pion-nucleus potential to an effective density-linear potential - generalized Seki-Masutani relations

Toshimitsu Yamazaki^{1*} and Satoru Hirenzaki²

¹*RI Beam Science Laboratory, RIKEN, 2-1 Hirosawa, Wako-shi, Saitama, 351-0198 Japan*

²*Department of Physics, Nara Women's University, Kita-Uoya Nishimachi, Nara 630-8506, Japan*
(October 29, 2002)

We have shown that a local-density-dependent term, $F[\rho(r)]\rho(r)$, of the pion-nucleus potential with a nuclear density, $\rho(r)$, can be transposed into a conventional density-linear term, $F(\rho_e)\rho(r)$, with an effective nuclear density, ρ_e , which is found to be close to $\sim 0.6\rho(0)$ for most π^- bound states. The presently found relations, *generalized Seki-Masutani relations*, for the density-quadratic term, the medium-modified isovector term and the double-scattering isoscalar term of the s-wave pion-nucleus interaction assure that the constant parameters in the conventional pion-nucleus potential are interpreted as being effective ones in the light of density-dependent effects.

1. Introduction

The so called “anomalous s-wave repulsion” in the pion-nucleus interaction has been a long-standing problem in pion physics (see, for instance, [1]). Many authors (see, for instance, [1,2]) have claimed the existence of a large extra repulsion arising from the density quadratic term, $\text{Re}B_0\rho(r)^2$, from global fits of pionic atom data to the Ericson-Ericson potential [3,4]:

$$V(r) = U_s(r) + U_p(r), \quad (1)$$

$$U_s(r) = -(2\pi/m_\pi) \times [\epsilon_1(b_0\rho(r) + b_1\Delta\rho(r)) + \epsilon_2 B_0\rho(r)^2],$$

$$\rho(r) = \rho_p(r) + \rho_n(r), \quad (2)$$

$$\Delta\rho(r) = \rho_n(r) - \rho_p(r), \quad (3)$$

where $U_s(r)$ and $U_p(r)$ are for s-wave and p-wave parts, respectively, and $\rho_p(r)$ and $\rho_n(r)$ are the proton and neutron density distributions, respectively, and $\epsilon_1 = 1 + m_\pi/M_N$ and $\epsilon_2 = 1 + m_\pi/2M_N$ (hereafter, we adopt units of m_π^{-1} , m_π^{-1} and m_π^{-4} for b_0 , b_1 and $\text{Re}B_0$, respectively). New light has been shed on this problem, namely, the possibility of an anomalous repulsion in the isovector part (b_1) as a unique signature of chiral symmetry restoration was pointed out [5,6]. In this respect, the deeply bound π^- states which were produced in heavy nuclei (in ^{207}Pb [7–10] and in ^{205}Pb [11,12]) are of particular importance, since the binding energies and widths of 1s π^- states depend nearly entirely on the s-wave potential (U_s) and, thus, only the 1s states in heavy nuclei ($N > Z$) provide unique information on the isovector part of the s-wave potential. In fact, the above conjecture has been proved by using new information from

the 1s π^- state in ^{205}Pb . The experiment indicates an enhanced isovector parameter, which is related to a decreased order parameter of chiral symmetry breaking in the nuclear medium, $b_1^{\text{free}}/b_1 = f_\pi^{*2}/f_\pi^2 = 0.78_{-0.09}^{+0.13}$ [12]. Recent global fits of pionic atom data by Friedman [13,14] have shown consistency with the above view.

Since the potential parameter b_1 is nuclear-density dependent in the theoretical context of Weise [5], a question can be raised as to the mutual relation in adopting the conventional Ericson-Ericson potential with constant coefficients (model C) and a modified local-density-dependent (model LDD) potential. In the present note we show that both methods are equally valid and are connected to each other, since a density-dependent term can be effectively linearized. Our procedure gives an answer to the following question: where do the bound pions probe the nuclear potential? This problem is closely related to the so-called *Seki-Masutani (SM) relation* [16], which emphasized a strong correlation between the density-linear isoscalar parameter (b_0) and the density-quadratic parameter ($\text{Re}B_0$). In a sense, we generalize the SM relation.

2. Where does a bound π^- probe the nuclear potential?

First, we examine which part (r) of the nuclear density, $\rho(r)$, is probed by a bound π^- . Two typical nuclei, ^{16}O and ^{208}Pb , are considered. The former possesses only shallow bound states, whereas the latter accommodates deeply bound states. The nuclear matter densities are assumed to take a 2-parameter Fermi distribution with a half-density radius (c_p and c_n) and a diffuseness parameter (z_p and z_n). The parameter values were chosen as follows: $c_p = 6.652$ fm, $c_n = 6.892$ fm, and $a_p = a_n = 0.5234$ fm for ^{208}Pb , and $c_p = c_n = 2.482$ fm and $a_p = a_n = 0.5234$ fm for ^{16}O . The central nucleon densities are: $\rho_0 = 0.150$ for ^{208}Pb and 0.172 for ^{16}O . We take known potential parameters for the p-wave parts [1] and a set of s-wave parameters ($b_0 = -0.028$, $b_1 = -0.12$, $\text{Re}B_0 = 0$ and $\text{Im}B_0 = 0.055$).

Figure 1 shows the π^- densities ($R_{nl}(r)^2$), the nuclear densities ($\rho(r)$) and the overlapping densities (namely, the nuclear densities probed by π^-), defined as

$$S(r) = \rho(r) |R_{nl}(r)|^2 r^2, \quad (4)$$

for ^{16}O (1s and 2p) and ^{208}Pb (1s, 2p and 3d). From these figures we notice that the overlapping density is

peaked at a radius slightly less than the half-density radius, *nearly independent* of the nucleus and the π^- quantum numbers. This means that the bound π^- effectively probes a fraction of the full nuclear density ($\rho_0 = \rho(0)$):

$$\rho_e \sim 0.60 \rho_0. \quad (5)$$

This is a key to intuitively understanding the following results of numerical calculations.

3. Seki-Masutani relation

Seki and Masutani [16] emphasized the presence of a strong correlation between b_0 and $\text{Re}B_0$ in describing pionic-atom binding energies and widths, which we call the *Seki-Masutani relation*. Toki *et al.* found the same correlation also exists for the deeply bound 1s and 2p states in ^{208}Pb [17,18]. Recently, this correlation has been revisited, and the following common relation has been established both theoretically and empirically [12]:

$$b_0^* \equiv b_0 + 0.215 \text{Re}B_0 = \text{constant}. \quad (6)$$

This means that the binding energies (and widths) are nearly unchanged by varying either b_0 or $\text{Re}B_0$ as long as these parameters are moved together to fulfill the above SM relation. In other words, none of b_0 and $\text{Re}B_0$ can be uniquely determined from given binding energies; only their combination, b_0^* , can be experimentally determined.

Actually, various global fits gave widely distributed values with large errors for these parameters. For instance, the best-fit values of $\{b_0, \text{Re}B_0\}$ are $\{0.024(15), -0.26(3)\}$ in KLTK90 [2] and $\{0.000(15), -0.14(7)\}$ in BFG97 [1]. With a different definition of b_0 , namely, $b_0^F = b_0 - \Delta b_0^{\text{DS}}$, in which the double-scattering term (Δb_0^{DS} , to be discussed later) is subtracted, Friedman obtained three different sets for $\{b_0^F, \text{Re}B_0\}$: $\{0.018(19), -0.14(4)\}$ (F02a) [13], $\{0.030(10), -0.21(4)\}$ (F02b) [14] and $\{0.020(10), -0.15(4)\}$ [14]. These widely split values are shown by large open circles with both vertical and horizontal error bars in Fig. 2. They appear to be almost meaningless, contrary to their claim that the parameter $\text{Re}B_0$ is well determined. On the other hands, a χ^2 fit with a gridding $\text{Re}B_0$ yields a series of points, as shown by closed squares (FG98 [15]) and by small open circles (F02a [13]), both of which indicate linear relations parallel to the SM correlation.

Recently, the value of b_0 with a gridding $\text{Re}B_0$ was precisely determined by χ^2 fits of the 1s π^- binding energies in six light symmetric nuclei (^{12}C , ^{14}N , ^{16}O , ^{20}Ne , ^{24}Mg and ^{28}Si) [12]. The obtained data sets, shown by closed circles in Fig. 2 (their sizes correspond to the error bars), prove the above SM relation (6) perfectly, yielding a precise value of

$$b_0^* = -0.028 \pm 0.001. \quad (7)$$

It is to be noted that the b_0^* composed of the widely distributed values of b_0 and $\text{Re}B_0$ in each of KLTK90,

BFG97, F02a and F02b would yield a value, close to $-0.030 \sim -0.034$ (after subtraction of the double scattering term in the case of F02a and F02b), which is not so much distributed as the individual values of b_0 and $\text{Re}B_0$. These authors seemed to be puzzled by the largeness of their $|\text{Re}B_0|$ values compared with $\text{Im}B_0 = 0.055$. On the other hand, they never took seriously the other discrepancy that their b_0 's (after subtraction of the double scattering term) take large positive values, meaning a strongly attractive b_0 in contrast to the nearly vanishing free πN value [19,20]. Such odd consequences from the global fits must be artifacts, arising from regarding b_0 and $\text{Re}B_0$ as independent uncorrelated parameters.

4. Seki-Masutani Ansatz and its generalization

Seki and Masutani [16] showed that the correlation between b_0 and $\text{Re}B_0$ can be understood by imposing an effective replacement of the expectation value of the density-quadratic term as,

$$< \rho(r)^2 > \rightarrow \rho_e < \rho(r) >, \quad (8)$$

which is expected to hold for any pionic atom state with a common value of ρ_e (*Seki-Masutani Ansatz*). They claimed an effective nuclear density $\rho_e \sim 0.50 \rho_0$. We understand that the localization of the overlapping densities, $S(r)$, near $r \sim r_c$, as shown in Fig. 1, is the key to justify the SM Ansatz.

We now examine how the Seki-Masutani Ansatz applies to various functional forms of the potential, $U_s(\rho(r))$. Let us add the following form to the real part of the s-wave pion-nucleus potential:

$$\Delta U_s(r) = F[\rho(r)]\rho(r), \quad (9)$$

where $F[\rho(r)]$ is a local-density-dependent (LDD) functional coefficient, representing an additional density-nonlinear potential involving a local density, $\rho(r)$. We solve the Klein-Gordon equation with LDD potentials and compare the numerical results with those without invoking LDD assumptions, thereby examining how the effects of the LDD potentials are expressed. Following the spirit of SM, we study how this effect is transposed into an effective coefficient,

$$F[\rho(r)]\rho(r) \rightarrow \overline{F}(\rho_e)\rho(r), \quad (10)$$

where $\overline{F}(\rho_e)$ is an effective constant parameter involving an effective nuclear density, ρ_e . Namely,

$$\overline{F}(\rho_e) = \frac{< F[\rho(r)]\rho(r) >}{< \rho(r) >}. \quad (11)$$

5. Polynomial case

Let us first consider the case of a density-polynomial term, which can be expressed as

$$\text{Re}U_s(r) = -(2\pi/m_\pi)\epsilon_1[b + B\rho(r) + D\rho(r)^2]\rho(r), \quad (12)$$

where $b = b_0 + (N - Z)/A \times b_1$. We performed numerical calculations of the 1s binding energies of ^{16}O and ^{208}Pb , and found strong correlations between B and b and between D and b , as expressed by

$$^{208}\text{Pb} : b + 0.25B + 0.031D = \text{const}, \quad (13)$$

$$^{16}\text{O} : b + 0.20B + 0.025D = \text{const}. \quad (14)$$

The first relation is the well known Seki-Masutani relation for the $B\rho(r)^2$ term, and the second relation is a new relation for the $D\rho(r)^3$ term.

The effect of the density-polynomial potential is asserted to be equivalent to the effect of an effective potential,

$$U_{SM}(r) = -(2\pi/m_\pi)\epsilon_1 \times [b + 2.826B\rho_e^{(n=2)} + 8.000D(\rho_e^{(n=3)})^2]\rho(r), \quad (15)$$

with ρ_e in units of fm^{-3} . We obtained the following effective densities:

$$\rho_e^{(n=2)} \sim 0.088 \sim 0.59\rho_0, \quad (16)$$

$$\rho_e^{(n=3)} \sim 0.062 \sim 0.41\rho_0, \quad (17)$$

for ^{208}Pb , and

$$\rho_e^{(n=2)} \sim 0.071 \sim 0.42\rho_0, \quad (18)$$

$$\rho_e^{(n=3)} \sim 0.056 \sim 0.33\rho_0, \quad (19)$$

for ^{16}O . The effective density for the $n = 2$ term is around $0.5\rho_0$, as given by SM, but that for the $n = 3$ term is smaller (the pion probes the $\rho(r)^3$ component more at a larger r).

The effective density does not depend much on the individual states, as we have found in the cases of ^{16}O and ^{208}Pb . We are therefore tempted to expect that the concept of the effective density applies to any functional form of $U(\rho(r))$.

6. Density-dependent b_1 parameter

Next, let us consider the case of an isovector s-wave potential which is expected to be enhanced due to a partial restoration of the chiral symmetry in the nuclear medium [5,6]. We assume the following potential according to Weise [5]:

$$b_1(r)\Delta\rho(r) = \frac{b_1^{free}}{1 - \alpha\rho(r)}\Delta\rho(r), \quad (20)$$

where α is a constant parameter representing the density-dependent effect. We examine whether or not the effect of the above LDD potential can be replaced by a conventional potential with a constant parameter involving an effective density ρ_e , as in

$$\bar{b}_1\Delta\rho(r) = \frac{b_1^{free}}{1 - \alpha\rho_e}\Delta\rho(r). \quad (21)$$

Numerical calculations of the binding energies yielded the dependence of B_{1s} on the parameter α , as shown in Fig. 3. On the other hand, the standard procedure assuming b_1 to be a constant gave another relation between B_{1s} and b_1 . The two relations are found to be nearly identical, if we take $b_1 = \bar{b}_1$ with

$$\rho_e \sim 0.090 \sim 0.60\rho_0. \quad (22)$$

The above finding indicates that the computational results starting from the LDD potential invoking a $\rho(r)$ -dependent b_1 parameter are equivalent to the results with a constant-parameter conventional potential; the LDD potential can be safely replaced by a density-linear conventional potential with a coefficient, eq.(21).

7. Double-scattering term

The isoscalar s-wave parameter in the free πN scattering, b_0^{free} , was determined experimentally to be nearly zero [19,20] in accordance with the Tomozawa-Weinberg theorem [21,22]. On the other hand, the double-scattering gives a major correction to the isoscalar term, as given by [3,4,23]

$$\Delta b_0^{\text{DS}} = -[(b_0^{free})^2 + 2b_1^2] < \frac{1}{r} > \quad (23)$$

$$= -[(b_0^{free})^2 + 2b_1^2] \frac{3}{2\pi} k_F(\rho), \quad (24)$$

where $k_F(\rho) = [(3\pi^2/2)\rho(r)]^{1/3}$. This correction is local-density dependent, since the correlation function, $< 1/r >$, is density-dependent and b_1 can also be density dependent.

Let us first examine how the above relation changes when we take into account the LDD effect on $< 1/r >$, while keeping b_1 as a constant parameter. We solved the KG equation with $\Delta b_0^{\text{DS}} = -\beta(\rho(r))^{1/3} b_1^2$ as an isoscalar part, and obtained the B_{1s} values in ^{16}O and ^{208}Pb as functions of β , which were compared with B_{1s} calculated with Δb_0^{DS} as a constant. Equating the two results in the form

$$\Delta \bar{b}_0^{\text{DS}} = -\beta(\rho_e)^{1/3} b_1^2, \quad (25)$$

we find the effective density, ρ_e , is nearly a constant, around $\sim 0.40\rho_0$ for ^{16}O and $\sim 0.54\rho_0$ for ^{208}Pb . Thus, the coefficient in the double-scattering correction is by a factor of $(\rho_e/\rho_0)^{1/3} \sim 0.73$ and ~ 0.81 (for ^{16}O and ^{208}Pb , respectively) changed over the conventional formula with $\rho_e = \rho_0$, yielding

$$\Delta b_0^{\text{DS}} \sim -1.52 b_1^2. \quad (26)$$

Next, we evaluated the full LDD double scattering effect with a density-dependent $b_1(\rho)$. The same form as

eq.(20) with α as a running parameter was adopted. The results were compared with those calculated with b_1 as a constant, as shown in Table I. The effective density, $\rho_e/\rho_0 \sim 0.60$, for ^{16}O and ^{208}Pb agrees well with that found for $b_1(\rho)$. This means that the $\bar{b}_1$ effectively converted from the isovector LDD potential term has the same effect as the density-dependent $b_1(\rho)$ in the double-scattering formula, eq.(26).

If we assert that the isoscalar potential strength b_0^* , eq.(7), is composed of the free b_0 (essentially zero), the double scattering term and $\text{Re}B_0$, we can deduce $\text{Re}B_0$ empirically as [12]

$$\text{Re}B_0 = \frac{b_0^* - \Delta b_0^{\text{DS}}}{0.215} \sim -0.038 \pm 0.025. \quad (27)$$

This value looks reasonable, when compared with $\text{Im}B_0 = 0.055$.

8. RIA correction

The present procedure can be applied to any kind of additional potential term. Friedman [13,14] takes into account the following correction

$$\Delta U_{\text{RIA}} = -\frac{2\pi}{m_\pi} \frac{3}{5} \frac{m}{M} d_0 [(1 + a\rho)^2 - 1] k_F(\rho)^2 \rho(r), \quad (28)$$

where $d_0 = -0.19 m_\pi^{-3}$ and $a = 2.7 \text{ fm}^3$. The effect of adding ΔU_{RIA} was calculated and was found to be equivalent to invoking an additional isoscalar term

$$\Delta b_0^{\text{RIA}} \sim -0.021. \quad (29)$$

This correction can be obtained also by setting $\rho(r) \rightarrow \rho_e \sim 0.5 \rho_0$.

9. Angle transformation term

Finally, we make some comments on the so called angle transformation term as an additional correction to the EE potential,

$$\Delta U_{\text{AT}} = -(2\pi/m_\pi)[(m_\pi/2M)\epsilon_1^{-1}\nabla^2[c_0\rho + c_1\Delta\rho] + (m_\pi/M)\epsilon_2^{-1}\nabla^2(C_0\rho_n\rho_p)], \quad (30)$$

which has been taken into account in the global fits of BFG97, FG98, F02a and F02b, but not in the present paper nor in Ref. [12]. This correction behaves like an s-wave potential, though it originates from the p-wave parameters. We examined how this correction is expressed in terms of the s-wave parameters. We found that the inclusion of this correction brings the following changes:

$$\Delta b_0^{\text{AT}} \approx +0.003, \quad (31)$$

for both ^{16}O and ^{208}Pb , and

$$\Delta \text{Im}B_0^{\text{AT}} \approx -0.008 \text{ for } ^{16}\text{O}, \quad (32)$$

$$\approx -0.014 \text{ for } ^{208}\text{Pb}. \quad (33)$$

Namely, this correction brings an attractive effect to the real part and a narrowing effect to the imaginary part. Thus, the best-fit parameters (b_0^*) with and without this correction have differences, as given above. A small difference in b_0^* between the above global fits and the present analysis is attributed to this correction. On the other hand, there is no sizable effect on b_1 : $\Delta b_1^{\text{AT}} \approx +0.003$.

10. Conclusion

We have shown that the Seki-Masutani relation between b_0 and $\text{Re}B_0$ holds for the 1s binding energies of light and heavy pionic atoms, both theoretically and empirically. If we admit this fact, the parameter for the isoscalar s-wave interaction is well represented by another parameter, b_0^* , as given in eq.(6,7), which takes care of the strong correlation between b_0 and $\text{Re}B_0$. The reason for this relation is understood from the fact that the overlapping density of any bound π^- with the nuclear density is peaked at the radius which is slightly smaller than the half-density radius; the nuclear density effectively probed by a bound π^- is around $0.6 \rho_0$.

We have shown that the Seki-Masutani Ansatz can be generalized for any functional form of the density-dependent potential: a LDD term, $F[\rho(r)]\rho(r)$, can be transposed into a corresponding term $F[\rho_e]\rho(r)$ with a constant parameter, where ρ_e is an effective parameter. Some important cases are summarized in Table II. In other words, we can say that the conventional potential form persists to be valid, because it can represent a LDD potential; the constant parameters in the conventional potential are regarded as being density-dependent parameters at $\rho \sim \rho_e \sim 0.60 \rho_0$. This also means that the observed enhancement of b_1 (interpreted as a reduction of the chiral order parameter) in ^{205}Pb [12], $R = b_1^{\text{free}}/b_1 = f_\pi^{*2}/f_\pi^2 = 0.78_{-0.09}^{+0.13}$, is for the effective nuclear density of $\rho_e \sim 0.60 \rho_0$, and thus, the reduction that would occur for the full nuclear density is expected to be $R \sim 0.63$.

The authors would like to thank Professors R.S. Hayano and P. Kienle for the stimulating discussions. This work is supported by the Grant-in-Aid of Monbukagakusho of Japan.

-
- [1] C. Batty, E. Friedman and A. Gal, Phys. Reports **287** (1997) 385, and references therein.
 - [2] J. Konijn, C.T.A.M. de Laat, A. Taal and J.H. Koch, Nucl. Phys. **A519** (1990) 773.
 - [3] M. Ericson and T.E.O. Ericson, Ann. Phys. **36** (1966) 323.
 - [4] T.E.O. Ericson and W. Weise, "Pions in Nuclei" (Clarendon Press, Oxford, 1988).

- [5] W. Weise, Acta Physica Polonica **31** (2000) 2715.
- [6] P. Kienle and T. Yamazaki, Phys. Lett. **B514** (2001) 1.
- [7] T. Yamazaki *et al.*, Z. Phys. **A355** (1996) 219.
- [8] T. Yamazaki *et al.*, Phys. Lett. **B418** (1998) 246.
- [9] H. Gilg *et al.*, Phys. Rev. **C62** (2000) 025201.
- [10] K. Itahashi *et al.*, Phys. Rev. **C62** (2000) 025202.
- [11] H. Geissel *et al.*, Phys. Rev. Lett. **88** (2002) 122301.
- [12] H. Geissel *et al.*, Phys. Lett. **B**, in press.
- [13] E. Friedman, Phys. Lett. **B524** (2002) 87.
- [14] E. Friedman, arXiv.nucl-th/0205054v2 (2002).
- [15] E. Friedman and A. Gal, Phys. Lett. **B432** (1998) 235.
- [16] R. Seki and K. Masutani, Phys. Rev. **C27** (1983) 2799.
- [17] H. Toki and T. Yamazaki, Phys. Lett. **B213** (1988) 129.
- [18] H. Toki, S. Hirenzaki, R.S. Hayano and T. Yamazaki, Nucl. Phys. **A501** (1989) 653.
- [19] H.-Ch. Schröder *et al.*, Phys. Lett. **B469** (1999) 25.
- [20] T.E.O. Ericson, B. Loiseau and A.W. Thomas, Nucl. Phys. **A684** (2001) 380.
- [21] Y. Tomozawa, Nuovo Cimento **A46** (1966) 707.
- [22] S. Weinberg, Phys. Rev. Lett. **17** (1966) 616.
- [23] N. Kaiser and W. Weise, Phys. Lett. **B512** (2001) 283.

TABLE I. Relation between the coefficient α representing the density-dependent $b_1(\rho(r)) = b_1^{\text{free}}/[1 - \alpha\rho(r)]$ and the equivalent isovector parameter $\bar{b}_1$ in giving the same 1s binding energies.

Nucleus	α	$1 - b_1^{\text{free}}/\bar{b}_1$	ρ_e	ρ_e/ρ_0
^{16}O	2	0.215	0.1075	0.63
	3	0.323	0.1077	0.63
	4	0.428	0.1070	0.63
^{208}Pb	2	0.185	0.0925	0.62
	3	0.277	0.0923	0.62
	4	0.370	0.0925	0.62

TABLE II. Summary of the effective densities, ρ_e/ρ_0 , obtained for various types of local-density-dependent functions, $F[\rho(r)]\rho(r)$.

$F[\rho]$	1s ^{16}O	1s ^{208}Pb	Remark
ρ	0.42	0.59	SM relation
$1/(1 - \alpha\rho)$		0.60	density-dependent b_1
$\rho^{1/3}$	0.40	0.54	effective Fermi momentum
$\rho^{1/3}/(1 - \alpha\rho)^2$	0.63	0.62	double-scattering term




FIG. 1. Overlapping probabilities (lower frame) of the π^- densities (upper frame) with the nucleon densities (middle frame) in typical pionic bound states: (Left) ^{16}O . (Right) ^{208}Pb . The vertical broken lines show the half-density proton radii and the vertical dash-dotted line is for the half-density neutron radius in ^{208}Pb .




FIG. 2. Seki-Masutani relation between b_0 and $\text{Re} B_0$. Best-fit values of b_0 versus $\text{Re} B_0$ as a gridding variable, obtained in χ^2 minimization using the 1s pionic atom data in 6 $N = Z$ nuclei, are shown by closed circles, whose sizes are equal to the fitting uncertainties. They lie on the SM lines: $b_0^* = b_0 + 0.215 \text{Re} B_0 = -0.0280 \pm 0.0010$. The $\{b_0, \text{Re} B_0\}$ parameters obtained from global fits of KLTK90 [2], BFG97 [1], F02a [13] and one of F02b [14] are shown by large open circles with both vertical and horizontal error bars. Also shown are $\{b_0, \text{Re} B_0\}$ sets from FG98 [15] (closed squares) and F02a [13] (small open circles), without conversion from b_0 to b_0^F in their convention.




FIG. 3. The 1s binding energy calculated in the LDD potential, eq.(20), as a function of α , compared with that in a conventional potential with b_1 as a constant parameter.